

DOCTORAL (PhD) DISSERTATION

University of Sopron

Roth Gyula Doctoral School of Forestry and Wildlife Management Sciences

**Understanding and Predicting the Potential Impacts of Climate Changes and
Anthropogenic Drivers on Savanna Ecosystems**

By:

Emad Hassan Elawad Yasin

Supervisor:

Prof. Dr. Kornel Czimer



Sopron

2026

1. Introduction

Climate change is a major global driver of ecosystem transformation, profoundly altering species distributions, vegetation structure, and ecological processes (IPCC, 2022; Yasin et al., 2024; Musa et al., 2024). Over the past century, global temperatures have increased by approximately 0.74 °C, with accelerated warming since the mid-1970s exceeding that of any comparable period in the last millennium (IPCC, 2022; Zohrabi et al., 2014; Babatolu & Akinnubi, 2013). Minimum temperatures are rising faster than maximum temperatures in many regions, increasing stress on species with narrow thermal tolerances (Hughes et al., 2003; Guan et al., 2015; Yasin et al., 2025). At the same time, atmospheric CO₂ concentrations have increased by nearly 100 ppm above pre-industrial levels, affecting plant physiology and ecosystem functioning (IPCC, 2014; IPCC, 2022; Coleine et al., 2024).

In addition to climatic changes, ecosystems are increasingly exposed to anthropogenic pressures such as agricultural expansion, deforestation, overgrazing, and habitat fragmentation, which reduce ecosystem resilience and adaptive capacity (Jump & Peñuelas, 2005; Musa & Sahoo, 2023; Abuelbashar et al., 2022). The interaction between climatic and human-induced stressors amplifies ecological vulnerability, with up to 30% of global species projected to face elevated extinction risk under combined pressures (Maclean & Wilson, 2011; Musa et al., 2024; Yasin & Mulyana, 2022).

Savanna ecosystems are particularly sensitive to these interacting drivers due to their structural heterogeneity, climatic variability, and ecological constraints (Colwell et al., 2008; Deutsch et al., 2008; Yasin et al., 2024). In Africa, savannas form extensive transitional landscapes between deserts and tropical forests and provide essential ecosystem services, including biodiversity conservation, carbon storage, soil stabilization, and livelihood support (Harrison & Jackson, 1958; Fahmi, 2017; Siddig et al., 2018; Musa & Sahoo, 2023; Yasin & Mulyana, 2022).

In Sudan, savanna ecosystems are structured by rainfall gradients, soil properties, fire regimes, and human activities, resulting in strong spatial variability in species composition (Fahmi, 2017; Musa et al., 2024; Gurashi et al., 2024). Drought-tolerant species dominate northern zones, while mesic southern regions support diverse woody species, including *Acacia seyal*, *Combretum hartmannianum*, and *Anogeissus leiocarpus* (Ibrahim & Osman, 2014; Yasin et al., 2024; Musa et al., 2024).

However, Sudanese savannas are increasingly undergoing degradation driven by rising temperatures, rainfall variability, and prolonged droughts, which affect tree growth, mortality, and regeneration (Fahmi, 2017; D'Odorico & Bhattachan, 2012; Yasin et al., 2025). These climatic stressors, combined with anthropogenic pressures such as deforestation and land-use expansion, lead to reduced woody cover, altered species composition, and declining ecosystem functionality (Smith et al., 2019; Musa et al., 2024; Coleine et al., 2024). Despite their importance, these ecosystems remain underrepresented in integrated climate impact assessments (Musa et al., 2024; Coleine et al., 2024).

A major limitation is the lack of spatially explicit and long-term integrated studies that simultaneously address climate variability, land-use change, vegetation dynamics, and species-level responses. Existing

research is often fragmented and rarely incorporates socio-economic and policy dimensions (Abuelbashar et al., 2022; Fahmi, 2017; Lu et al., 2023; Musa et al., 2024; Hemida et al., 2023).

Recent advances in remote sensing and machine learning provide powerful tools to overcome these limitations. Multi-temporal satellite data enable large-scale monitoring of vegetation dynamics, while algorithms such as Random Forest (RF), Support Vector Machine (SVM), and Classification and Regression Trees (CART) support robust modeling of complex ecological relationships (Gorelick et al., 2017; Huete et al., 2002; Belgiu & Drăguț, 2016; Pal, 2005; Raczko & Zagajewski, 2017). In addition, species distribution models (SDMs) enable prediction of species responses to future climate scenarios and identification of climate refugia (Moss et al., 2010; Ludwig et al., 2013; Yasin et al., 2024; Coleine et al., 2024).

However, the application of these approaches in Sudan remains limited, with few studies integrating climate analysis, land-use dynamics, ecological data, machine learning, and species distribution modeling within a unified framework.

Therefore, this study develops an integrated analytical framework to assess the combined impacts of climatic variability and anthropogenic pressures on Sudanese savanna ecosystems. This framework provides a robust scientific basis for conservation planning, adaptive management, and climate change mitigation strategies.

1.1. General Objective

The general objective of this study is to understand and predict the combined impacts of climatic variability and anthropogenic pressures on savanna ecosystems in Sudan by integrating climate analysis, land use/land cover dynamics, field-based ecological assessment, and geospatial machine learning approaches.

This study addresses the following specific objectives:

1. To analyze the long-term climatic variability, trends, and drought patterns in savanna woodlands of southeastern Sudan.
2. To assess the spatio-temporal land use/land cover (LULC) changes and their interactions with climatic variability.
3. To evaluate the changes in woody species composition, diversity, and regeneration in response to climatic variability and anthropogenic pressures.
4. To map the dominant tree species using multi-source remote sensing data and machine learning approaches.
5. To model and predict the current and future potential distribution of *Acacia seyal* (Delile) under different climate change scenarios.

2. Literature Review

The literature review includes scientific studies related to the main themes addressed in this thesis, including Sudanese savanna ecosystems, climate variability and drought, land use/land cover (LULC)

dynamics, vegetation monitoring and modeling, remote sensing and Machine Learning applications, species distribution modeling, and sustainable management and climate adaptation strategies.

3. Research Methodology

3.1. Study Area

The study was conducted in Sudan, with a focus on Blue Nile State and the Elnour Natural Forest Reserve (ENFR). Sudan exhibits strong climatic and ecological gradients, ranging from hyper-arid conditions in the north to semi-humid savanna ecosystems in the south, which strongly influence vegetation distribution and ecosystem dynamics. Blue Nile State, located in southeastern Sudan, represents a transitional zone characterized by a tropical savanna climate with a distinct wet season (April–November) and dry season (December–March), annual rainfall ranging from 300 to 700 mm, and generally high temperatures. ENFR, situated within Blue Nile State, is a low-rainfall woodland savanna established for biodiversity conservation, covering approximately 4,667 ha. The reserve supports diverse woody vegetation dominated by species such as *Acacia seyal* (Delile), *Combretum hartmannianum*, and *Anogeissus leiocarpus*. The area is influenced by surrounding communities through grazing and resource use, contributing to ongoing vegetation change and degradation processes.

3.2 Data Collection

- This study integrated remote sensing data, field observations, climatic datasets, and species occurrence records to assess the impacts of climatic variability and anthropogenic pressures on savanna ecosystems.
- Long-term climate data, including rainfall and temperature, were obtained from the Climatic Research Unit (CRU) dataset and complemented with station observations from the Sudan Meteorological Authority to ensure data reliability and regional consistency. Future climate projections were derived from CMIP6 datasets for scenario-based analysis.
- Multi-temporal satellite imagery from Landsat 5 Thematic Mapper (2008), Landsat 8 Operational Land Imager (2013), and Sentinel-2 Multispectral Instrument (2018 and 2021) was acquired through Google Earth Engine for different reference years to analyze land use/land cover dynamics and vegetation patterns.
- Field data were collected from 229 permanent sample plots established in the Elnour Natural Forest Reserve, where measurements of species composition, tree diameter, height, and regeneration status were recorded during surveys conducted in 2008 and revisited in 2013, 2018, and 2021.
- In addition, environmental variables, including bioclimatic data, soil properties, and elevation, were obtained from global datasets such as WorldClim and SoilGrids to support species distribution modeling. Occurrence records of *Acacia seyal* were compiled from field surveys,

national forest inventories, biodiversity databases, and published studies. These records were standardized and filtered to ensure spatial accuracy and reduce sampling bias.

3.3 Data Analysis

- Data processing and analysis were carried out using a combination of cloud-based platforms, geographic information systems, statistical software, and modeling tools. Climate data were analyzed using Google Earth Engine, R, Statistica, and Microsoft Excel to assess variability, trends, and drought characteristics through the application of the Mann–Kendall test, Sen’s slope estimator, and the Standardized Precipitation Evapotranspiration Index (SPEI).
- Satellite imagery was preprocessed and analyzed in Google Earth Engine to generate land use/land cover maps through supervised classification techniques, primarily using the Random Forest algorithm. Change detection and transition analysis were conducted to quantify spatio-temporal landscape dynamics, while accuracy assessment was performed using standard classification metrics.
- Field-based ecological data were analyzed using R, Statistica, and Excel to evaluate species composition, diversity, stand structure, and regeneration status through the application of standard phytosociological and biodiversity indices.
- For tree species mapping, Machine Learning algorithms, including Random Forest, Support Vector Machine, Classification and Regression Trees, and ensemble approaches, were implemented within Google Earth Engine to classify dominant species and assess model performance.
- Species distribution modeling was conducted using the MaxEnt model to predict the current and future distribution of *Acacia seyal* (Deliel). Environmental variables were screened for multicollinearity, and model performance was evaluated using AUC and TSS metrics. Future projections were generated under different climate change scenarios (SSP2–4.5 and SSP5–8.5), and habitat suitability maps were produced to identify potential climate refugia and vulnerable areas.

4. New Scientific Results

This dissertation presents original scientific contributions to understanding land degradation processes, biodiversity dynamics, forest structural change, and climate-driven species distribution in Sudanese savanna ecosystems. These findings are derived from an integrated multi-scale framework combining climate analysis, remote sensing, field observations, and modeling approaches, and are summarized as follows:

4.1. Climate–Drought Mechanism

I demonstrate that drought dynamics in Sudanese savanna ecosystems are increasingly influenced by temperature and rainfall variability rather than total precipitation alone. Using long-term climate data

(1950–2024) and multi-scale SPEI analysis, I show that rising temperatures amplify atmospheric evaporative demand, intensifying moisture stress even under stable or slightly increasing rainfall. The results provide strong empirical evidence of a transition from precipitation-dominated to energy-influenced drought processes. I further establish a process-based link between climate variability and ecosystem response, identifying temperature-driven moisture stress as a key factor contributing to vegetation degradation, reduced regeneration, and structural change.

Related publication:

Yasin, E. H. E., Yagoub, Y. E., & Czimber, K. (2026). Long-Term Climate Variability, Trends, and Future Drought Projections in Blue Nile State, Sudan (1901–2100). *Journal of Hydrology* (***Submitted***).

4.2. Semi-Bareland as a Degradation Pathway

I demonstrate that semi-bareland has become the dominant and persistent land-use/land-cover class in Elnour Natural Forest Reserve (ENFR) between 1995 and 2021. Based on multi-temporal LULC analysis and transition dynamics, the results indicate that semi-bareland represents a major and sustained degradation pathway in the study area, reflecting a progressive transition from forested to more degraded states under continued disturbance. While the findings strongly support its persistence and dominance, its characterization as a fully stable alternative state would require additional long-term ecological and process-based evidence.

4.3. Drivers of Forest Degradation

I demonstrate, based on integrated field observations, land-use/land-cover analysis, and multiple lines of evidence, that anthropogenic activities are the primary drivers of forest degradation in ENFR. Illegal logging, charcoal production, and overgrazing emerge as the dominant pressures contributing to forest decline, while governance and socio-economic factors further exacerbate degradation. Climatic variability acts primarily as a secondary amplifier, intensifying the effects of human-induced disturbances rather than functioning as the principal driver.

Related publication for (4.2 and 4.3):

Yasin, E. H. E., Koren, M., & Czimber, K. (2026). Spatio-temporal assessment and future projection of land cover dynamics in savanna woodlands of Sudan using Machine Learning and CA–ANN modeling. *Remote Sensing*, 18(7), 1086; <https://doi.org/10.3390/rs18071086>

4.4. Biodiversity Loss and Regeneration Failure

I demonstrate a substantial decline in woody species richness from 35 to 19 species between 2008 and 2021, accompanied by widespread regeneration failure and the absence of successful recruitment in several species. The results indicate a severe limitation in regeneration capacity, which represents a key process driving biodiversity loss and reduced ecosystem resilience.

4.5. Structural Simplification of Forests

I demonstrate that forest structure is increasingly dominated by small diameter classes (exceeding 43%), while large trees (below 12%) have declined significantly. The results indicate a progressive structural simplification of the forest, characterized by reduced representation of mature trees and increased dominance of smaller size classes. This pattern indicates that selective pressures, including the removal of mature trees and other disturbances, are associated with disrupted cohort development and limited canopy recovery despite ongoing recruitment, revealing a decoupling between regeneration and structural recovery as a key factor contributing to persistent forest degradation.

Related publication for (4.4 and 4.5):

Yasin, E. H. E., Gibreel, H. H., Siddig, A. A. H., Musa, F. I., Sahoo, U. K., & Czimber, K. (2026). Spatiotemporal changes in woody species composition, diversity, and regeneration in a rapidly degrading savanna woodland in Blue Nile State, Sudan. *Frontiers in Forests and Global Change (Submitted)*.

4.6. Machine Learning Robustness in Species Mapping

I demonstrate, based on multi-temporal and multi-sensor classification results, that Random Forest (RF) and Support Vector Machine (SVM) consistently achieve high and stable performance under heterogeneous dryland conditions. The results show that reliable species mapping in Sudanese savanna ecosystems depends on algorithm robustness to class imbalance, mixed-pixel effects, and landscape heterogeneity, rather than on overall accuracy alone. This provides a structured evaluation of classification algorithms under multi-temporal, multi-sensor, and ecologically complex dryland conditions.

Related publication:

Yasin, E. H. E., Koreň, M., & Czimber, K. (2026). Machine Learning-Based Mapping of Dominant Tree Species in Dryland Forests Using Multi-Temporal and Multi-Source Data. *Remote Sensing*, 18(8), 1185. <https://doi.org/10.3390/rs18081185>

4.7. Climate-Driven Redistribution of *Acacia seyal*

I demonstrate, based on species distribution modelling, that the potential distribution of *Acacia seyal* is primarily influenced by isothermality and precipitation during the warmest quarter. The results indicate that projected climate change may lead to habitat contraction in northern regions and expansion in southern and eastern regions, identifying these areas as potential climatic refugia. These projections represent climatic suitability; their realization may be influenced by land use, disturbance regimes, and dispersal constraints, which are not explicitly included in the model.

Related publication:

Yasin, E. H. E., Siddig, A. A. H., Koren, M., & Czimer, K. (2026). Modeling the effects of climate change on the current and future potential distribution of *Acacia seyal* (Delile) P.J.H. Hurter in Sudan using a machine learning approach. *Modeling Earth Systems and Environment* (*Submitted*).

References

- Abuelbasha, A. I., Ahmed, D. A. M. D. A., Siddig, A. A. H., Yagoub, Y. E., & Gibreel, H. H. (2022). Analysis of Composition and Diversity of Natural Regeneration of Woody Species in Jebel El Gerrie Dry Land Forest East of Blue Nile State, Sudan. *Journal of forest and environmental science*, 38(2), 90-101. <https://doi.org/10.7747/JFES.2022.38.2.90>
- Babatolu, J. S., & Akinnubi, R. T. (2013). Surface temperature anomalies in the river Niger basin development authority areas, Nigeria. *Atmospheric and Climate Sciences*, 3(4), 532-537. <https://doi.org/10.4236/acs.2013.34056>
- Belgiu, M., & Drăguț, L. (2016). Random forest in remote sensing: A review of applications and future directions. *ISPRS journal of photogrammetry and remote sensing*, 114, 24-31. <https://doi.org/10.1016/j.isprsjprs.2016.01.011>
- Coleine, C., Delgado-Baquerizo, M., DiRuggiero, J., Guirado, E., Harfouche, A. L., Perez-Fernandez, C., ... & Egidi, E. (2024). Dryland microbiomes reveal community adaptations to desertification and climate change. *The ISME journal*, 18(1), wrae056. <https://doi.org/10.1093/ismejo/wrae056>
- Coleine, C., Delgado-Baquerizo, M., DiRuggiero, J., Guirado, E., Harfouche, A. L., Perez-Fernandez, C., ... & Egidi, E. (2024). Dryland microbiomes reveal community adaptations to desertification and climate change. *The ISME journal*, 18(1), wrae056. <https://doi.org/10.1093/ismejo/wrae056>
- Colwell, R. K., Brehm, G., Cardelús, C. L., Gilman, A. C., & Longino, J. T. (2008). Global warming, elevational range shifts, and lowland biotic attrition in the wet tropics. *science*, 322(5899), 258-261. <https://doi.org/10.1126/science.1162547>
- Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P. R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, 105(18), 6668-6672. <https://doi.org/10.1073/pnas.0709472105>
- D'Odorico, P., & Bhattachan, A. (2012). Hydrologic variability in dryland regions: impacts on ecosystem dynamics and food security. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1606), 3145-3157. <https://doi.org/10.1098/rstb.2012.0016>
- Fahmi, M. K. M. (2017). Climate, trees and agricultural practices: Implications for food Security in the semi-arid zone of Sudan. Univ. Helsinki Tropic. Forest. Dep., Helsinki.

- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote sensing of Environment*, 202, 18-27. <https://doi.org/10.1016/j.rse.2017.06.031>
- Guan, Y., Zhang, X., Zheng, F., & Wang, B. (2015). Trends and variability of daily temperature extremes during 1960–2012 in the Yangtze River Basin, China. *Global and Planetary Change*, 124, 79-94. <https://doi.org/10.1016/j.gloplacha.2014.11.008>
- Gurashi, N. A., Yasin, E. H. E., & Czimber, K. (2024). Changes in Structure, Tree Species Composition, and Diversity of the Abu Geili Riverine Forest Reserve, Sinnar State, Sudan. *Acta Silvatica Et Lignaria Hungarica: An International Journal In Forest, Wood and Environmental Sciences*, 20(2), 55-70. <https://doi.org/10.37045/aslh-2024-0004>
- Harrison, M. N., & Jackson, J. K. (1958) Ecological classification of the vegetation of the Sudan. *Forests Bulletin, Forest Department, Ministry of Agriculture, Khartoum, Sudan* 1958(2), 45.
- Hemida, M., Yasin, E. H. E., Kheiry, M. A., Hammad, Z. M., & Vityi, A. (2023). Assessment of Taungya agroforestry system in dryland forests rehabilitation in Sudan. *Journal of Degraded and Mining Lands Management*, 10, 4495–4507. <https://doi.org/10.15243/jdmlm.2023.103.4495>
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote sensing of environment*, 83(1-2), 195-213. [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2)
- Hughes, T. P., Baird, A. H., Bellwood, D. R., Card, M., Connolly, S. R., Folke, C., ... & Roughgarden, J. (2003). Climate change, human impacts, and the resilience of coral reefs. *science*, 301(5635), 929-933. <https://doi.org/10.1126/science.1085046>
- Ibrahim, E. M. I., & Osman, E. H. (2014). Diameter at Breast Height-Crown Width Prediction Models for *Anogeissus Leiocarpus* (DC.) Guill & Perr and *Combretum Hartmannianum* Schweinf. *Journal of Forest Products & Industries*, 3(4), 191-197.
- IPCC. (2014). Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
- IPCC. (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. In H. O. Pörtner, D. C. Roberts, M. Tignor, E. S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, & B. Rama (Eds.). Cambridge University Press.
- Jump, A. S., & Peñuelas, J. (2005). Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology letters*, 8(9), 1010-1020. <https://doi.org/10.1111/j.1461-0248.2005.00796.x>

- Lu, Y., Zhang, B., Zhang, M., Jie, M., Guo, S., & Wang, Y. (2023). Relict plants are better able to adapt to climate change: evidence from desert shrub communities. *Plants*, 12(23), 4065. <https://doi.org/10.3390/plants12234065>
- Ludwig, F., Supit, I., Franssen, W., & Biemans, H. (2013). *Climate change impacts on hydrological and meteorological extremes: Executive summary of COMBINE project*.
- Maclean, I. M., & Wilson, R. J. (2011). Recent ecological responses to climate change support predictions of high extinction risk. *Proceedings of the National Academy of Sciences*, 108(30), 12337-12342. <https://doi.org/10.1073/pnas.1017352108>
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., ... & Wilbanks, T. J. (2010). The next generation of scenarios for climate change research and assessment. *Nature*, 463(7282), 747-756. <https://doi.org/10.1038/nature08823>
- Musa, F. I., & Sahoo, U. K. (2023). Role of sustainable forest management in poverty reduction and livelihood improvement in Sudan: A review. *International Journal of Ecology and Environmental Sciences*, 49, 449-456. <https://doi.org/10.55863/ijees.2023.3036>
- Musa, F. I., Mohammed, M. H., Fragallah, S. D., Adam, H. E., & Sahoo, U. K. (2024). Current status of tree species diversity at Abu Gadaf Natural Forest Reserve, Blue Nile Region—Sudan. *Vegetos*, 37(5), 1760-1771. <https://doi.org/10.1007/s42535-024-00931-2>
- Pal, M. (2005). Random forest classifier for remote sensing classification. *International journal of remote sensing*, 26(1), 217-222. <https://doi.org/10.1080/01431160412331269698>
- Raczko, E., & Zagajewski, B. (2017). Comparison of support vector machine, random forest and neural network classifiers for tree species classification on airborne hyperspectral APEX images. *European Journal of Remote Sensing*, 50(1), 144-154. <https://doi.org/10.1080/22797254.2017.1299557>
- Siddig, A. A., Magid, T. D. A., EL-Nasry, H. M., Hano, A. I., & Mohammed, A. A. (2018). Biodiversity in Sudan. In *Global biodiversity*. Apple Academic Press. pp. 273-294. <https://doi.org/10.1201/9780429469800-10>
- Smith, W. K., Dannenberg, M. P., Yan, D., Herrmann, S., Barnes, M. L., Barron-Gafford, G. A., ... & Yang, J. (2019). Remote sensing of dryland ecosystem structure and function: Progress, challenges, and opportunities. *Remote Sensing of Environment*, 233, 111401. <https://doi.org/10.1016/j.rse.2019.111401>
- Yasin, E. H. E., & Mulyana, B. (2022). Spatial distribution of tree species composition and carbon stock in Tozi tropical dry forest, Sinnar State, Sudan. *Biodiversitas*, 23, 2359–2368. <https://doi.org/10.13057/biodiv/d230513>
- Yasin, E. H. E., Siddig, A. A. H., & Kornel, C. (2024). Forests at the Crossroads: Biodiversity Conservation in the Era of Climate Change. In S. N. Kulshreshtha (Ed.). *Sustainable Forest Management - Surpassing Climate Change and Land Degradation*. *Intech Open*. <https://doi.org/10.5772/intechopen.1004224>

- Yasin, E. H., Kamil, O. H., & Mulyana, B. (2022). Multi-temporal satellite images analysis for assessing and mapping deforestation in Um Hataba Forest, South Kordofan, Sudan. *Jurnal Sylva Indonesiana*, 5, 81–92. <https://doi.org/10.32734/jsi.v5i01.7504>
- Yasin, E. H., Siddig, A. A., Diab, E. E., & Czimber, K. (2025). Evaluating the Efficiency of Two Ecological Indices in Monitoring Forest Degradation in the Drylands of Sudan. *Remote Sensing*, 17(13), 2298. <https://doi.org/10.3390/rs17132298>
- Zohrabi, N., Bavani, A. M., Goodarzi, E., & Eslamian, S. (2014). Attribution of temperature and precipitation changes to greenhouse gases in northwest Iran. *Quaternary International*, 345, 130-137. <https://doi.org/10.1016/j.quaint.2014.01.026>

Publications list

- Yasin, E. H. E., Yagoub, Y. E., & Czimber, K. (2026).** Long-Term Climate Variability, Trends, and Future Drought Projections in Blue Nile State, Sudan (1901–2100). *Journal of Hydrology (Submitted)*.
- Yasin, E. H. E., Koren, M., & Czimber, K. (2026).** Spatio-temporal assessment and future projection of land cover dynamics in savanna woodlands of Sudan using Machine Learning and CA–ANN modeling. *Remote Sensing*, 18(7), 1086; <https://doi.org/10.3390/rs18071086>.
- Yasin, E. H. E., Gibreel, H. H., Siddig, A. A. H., Musa, F. I., Sahoo, U. K., & Czimber, K. (2026).** Spatiotemporal changes in woody species composition, diversity, and regeneration in a rapidly degrading savanna woodland in Blue Nile State, Sudan. *Frontiers in Forests and Global Change (Submitted)*.
- Yasin, E. H. E., Koren, M., & Czimber, K. (2026).** Machine Learning-Based Mapping of Dominant Tree Species in Dryland Forests Using Multi-Temporal and Multi-Source Data. *Remote Sensing*, 18(8), 1185. <https://doi.org/10.3390/rs18081185>
- Yasin, E. H. E., Siddig, A. A. H., Koren, M., & Czimber, K. (2026).** Modeling the effects of climate change on the current and future potential distribution of *Acacia seyal* (Delile) P.J.H. Hurter in Sudan using a machine learning approach. *Modeling Earth Systems and Environment (Submitted)*.
- Yasin, E. H. E., Siddig, A. A. H., Diab, E. E., & Czimber, K. (2025).** Evaluating the efficiency of two ecological indices in monitoring forest degradation in the drylands of Sudan. *Remote Sensing*, 17(13), 2298. <https://doi.org/10.3390/rs17132298>
- Yasin, E. H. E., Sharif, M. M., Yahia, M. Y., Othman, A. Y., Ibrahim, A. O., Kheiry, M. A., & Musa, M. (2024).** Crop type classification and spatial mapping in River Nile and Northern State, Sudan, using Sentinel-2 satellite data and field observation. *Journal of Degraded and Mining Lands Management*, 11(3), 5997–6007. <https://doi.org/10.15243/jdmlm.2024.113.5997>

- Yasin, E. H. E.,** Mulyana, B. (2022). Spatial distribution of tree species composition and carbon stock in Tozi tropical dry forest, Sinnar State, Sudan. *Biodiversitas*, 23, 2359–2368. <https://doi.org/10.13057/biodiv/d230513>
- Yasin, E. H. E.,** Kamil, O., & Mulyana, B. (2022). Multi-temporal satellite images analysis for assessing and mapping deforestation in Um Hataba Forest, South Kordofan, Sudan. *Journal of Sylva Indonesiana*, 5(1), 81–92. <https://doi.org/10.32734/jsi.v5i01.7504>
- Mohamed, A. A. A., Musa, F. I., Lalruatkimi, C., **Yasin, E. H. E.,** Thawmmawii, L., Alfadol, A. I. A., Elyas, A. N., Khanna, N., Ashish, P., Tripathi, O. P., Abdelmagid, T. D., Sahoo, U. K., & Abushambo, A. A. K. (2026). Satellite-based drought assessment in major Sudan's mechanized rain-fed agriculture. *Scientific Reports*. <https://doi.org/10.1038/s41598-026-49618-5>
- Elgaber, A. K. A., **Yasin, E. H. E.,** Czimber, K., Osman, A. G., & Elsheikh, E. A. (2026). Assessment of the Effect of Atrazine on Soil Algal Flora in Sudan. *Applied and Environmental Soil Science*, 2026(1), 5594923. <https://doi.org/10.1155/aess/5594923>.
- Arbain, M. K. A., Khiry, M. A., Koren, M., & **Yasin, E. H. E.** (2026). Multi-temporal Landsat images analysis for forest degradation appraisal (1988–2018) in Umfakarin Natural Forest Reserve, Sudan. *International Journal of Geoinformatics*, 22(3), 99–116. <https://doi.org/10.52939/ijg.v22i3.4869>
- Hasoba, A. M. M., Yasin, E. H. E., Osman, M. B. O., & Czimber, K. (2025). Monitoring ecosystem dynamics using machine learning: Random forest-based LULC analysis in Dinder Biosphere Reserve, Sudan. *Engineering Proceedings*, 94(1), 2. <https://doi.org/10.3390/engproc2025094002>
- Elgaber, A. K. A., **Yasin, E. H. E.,** Mustafa, M., Zubairu, A. M., Czimber, K., Osman, A. G., & Elsheikh, E. A. E. (2025). Impact of NPK, plant residue, soil type, and temperature on the half-life of atrazine herbicide. *Nitrogen*, 6(3), 79. <https://doi.org/10.3390/nitrogen6030079>
- Mohammed, E. M. I., **Yasin, E. H. E.,** & Mulyana, B. (2025). Analysis of species richness variability, spatial distribution, and carbon stock in Okalma Natural Forest Reserve, Sudan. *Anthropocene Science*, 4(1), 30–41. <https://doi.org/10.1007/s44177-025-00091-1>
- Musa, F. I., Sahoo, U. K., Mohamed, A. A. A., Alfadol, A. I. A., Chettri, M., Ibrahim, E. M. I. M., Abdel Magid, T. D., **Yasin, E. H. E.,** Mohammed, M. H., Eltahir, M. E. S., & Azrag, M. A. M. (2025). Seed dispersal: Methods, agents, global research trends and networks. *Phytomorphology*, 75, 121–141. <https://doi.org/10.56024/ispm.75.3ampersand4.2025/121-141>
- Gurashi, N. A., **Yasin, E. H. E.,** & Czimber, K. (2024). Changes in structure, tree species composition, and diversity of the Abu Geili Riverine Forest Reserve, Sinnar State, Sudan. *Acta Silvatica & Lignaria Hungarica*, 20(2), 55–70. <https://doi.org/10.37045/aslh-2024-0004>

- Gurashi, N. A., **Yasin, E. H. E.**, & Czimber, K. (2024). Assessment of tree species availability based on sawmilling and timber markets survey in Sinnar State, Sudan. *Acta Silvatica & Lignaria Hungarica*, 20(1), 39–51. <https://doi.org/10.37045/aslh-2024-0003>
- Osman, M. B., & **Yasin, E. H. E.** (2024). Fostering environmental and resources management in Sudan through geo-information systems: A prospective approach for sustainability. *Journal of Degraded and Mining Lands Management*, 11(3), 5647–5657. <https://doi.org/10.15243/jdmlm.2024.113.5647>
- Elzubair, A. E., **Yasin, E. H. E.**, Alla, D. A. M. D., & Sanjak, E. (2024). Assessment of the role of Gum Arabic Producers' Associations (GAPAs) in rehabilitation of gum Arabic belt: Case of North and West Kordofan States, Sudan. *International Journal of Conservation Science*, 15(1), 461–478. <https://doi.org/10.36868/IJCS.2024.01.06>
- Hemida, M., **Yasin, E. H. E.**, Kheiry, M. A., Hammad, Z. M., & Vityi, A. (2023). Assessment of Taungya agroforestry system in dryland forests rehabilitation in Sudan. *Journal of Degraded and Mining Lands Management*, 10(3), 4495–4507. <https://doi.org/10.15243/jdmlm.2023.103.4495>
- Matatula, J., Wirabuana, P. Y. A. P., **Yasin, E. H. E.**, & Mulyana, B. (2023). Species composition and carbon stock of rehabilitated mangrove forest in Kupang District, East Nusa Tenggara, Indonesia. *Environmental Research Engineering and Management*, 79(3), 24–34. <https://doi.org/10.5755/j01.ereem.79.3.33123>
- Ahmed, H. O., Kheiry, M. A., & **Yasin, E. H. E.** (2023). Land-use land-cover change detection using geospatial techniques in Zalingei, Central Darfur, Sudan. *Journal of Information Systems and Informatics*, 5(4), 1676–1690. <https://doi.org/10.51519/journalisi.v5i4.624>
- Koko, S. A., Siddig, A. A., **Yasin, E. H. E.**, Gadallah, N. A., Adam, M., Hasoba, A. M., & Mohammed, A. A. (2023). Do observations during patrolling trips detect changes in wildlife presence and diversity in national parks? *Open Journal of Ecology*, 13(11), 794–805. <https://doi.org/10.4236/oje.2023.1311049>
- Elzaki, I. A. E., Siddig, A. A. H., **Yasin, E. H. E.**, Attaelmnan, A. A., Gadallah, N. A. H., Hasoba, A. M. M., Nasreldeen, M. A., Yagoub, Y. E., & Czimber, K. (2023). Effect of simulated drought and rainfall fluctuation on seedling growth of two savannah tree species in Sudan: An experimental exploration. *Acta Silvatica & Lignaria Hungarica*, 19(1), 37–50. <https://doi.org/10.37045/aslh-2023-0003>
- Purwanto, R. H., Mulyana, B., Satria, R. A., **Yasin, E. H. E.**, Putra, I. S. R., & Putra, A. D. (2022). Spatial distribution of mangrove vegetation species, salinity, and mud thickness in mangrove forest in Pangarengan, Cirebon, Indonesia. *Biodiversitas*, 23(3), 1384–1392. <https://doi.org/10.13057/biodiv/d230324>
- Eisawi, K. A., Subedi, I. P., **Yasin, E. H. E.**, Yode, C., & He, H. (2022). Variation in the rates of biomass removal by soil macro-fauna in different land uses at Rashad, South Kordofan,

- Sudan. Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa", 65(2), 173–197. <https://doi.org/10.3897/travaux.65.e95270>
- Younis, A. Y. I., Jonathan, B. N., Togun, A. O., **Yasin, E. H. E.**, Omer, S. O., Hamad, G. E. A. I., & Aïssata, S. D. (2022). Assessment of land use and land cover change on Gum Talha (*Acacia seyal* var. *seyal*) forest in Bahar Alarab, Sudan. *Biodiversitas*, 23(9), 4549–4560. <https://doi.org/10.13057/biodiv/d230922>
- Purwanto, R. H., Mulyana, B., Satria, R. A., **Yasin, E. H. E.**, Raditya Putra, I. S., & Putra, A. D. (2022). Spatial distribution of mangrove vegetation species, salinity, and mud thickness in mangrove forest in Pangarengan, Cirebon, Indonesia. *Biodiversitas: Journal of Biological Diversity*, 23(3), 1383–1391. <https://smujo.id/biodiv/article/view/10516/5585>
- Eisawi, K. A., Gibreel, H. H., Hong, H., Shaheen, T., Abdalla, O. M., & **Yasin, E. H. E.** (2022). Ethnobotanical study of medicinal trees and shrubs from the Rashad District of Southern Kordofan, Sudan. *Egyptian Journal of Botany*, 62(2), 337–357.
- Eisawi, K. A., He, H., Shaheen, T., & **Yasin, E. H. E.** (2021). Assessment of tree diversity and abundance in Rashad Natural Reserved Forest, South Kordofan, Sudan. *Open Journal of Forest*, 11, 37–46. <https://doi.org/10.4236/ojf.2021.111003>

Book chapters

- Yasin, E. H. E.**, Czimber, K., Gadallah, N. A. H., Mohammed, S. M. F., & Yagoub, Y. E. (2025). Challenges and future opportunities of geospatial modeling for natural resource assessment. In S. Das et al. (Eds.), *Application of Geospatial Technology and Modelling on Natural Resources Management. Advances in Geographic Information Science*. Springer, Cham. https://doi.org/10.1007/978-3-031-84698-4_22
- Yasin, E. H. E.**, Czimber, K., Gadallah, N. A. H., Mohammed, S. M. F., & Yagoub, Y. E. (2025). Geospatial techniques for mapping and monitoring land use land cover change. In S. Das et al. (Eds.), *Application of Geospatial Technology and Modelling on Natural Resources Management. Advances in Geographic Information Science*. Springer, Cham. https://doi.org/10.1007/978-3-031-84698-4_14
- Yasin, E. H. E.**, Czimber, K., Gadallah, N. A. H., Mohammed, S. M. F., & Yagoub, Y. E. (2025). Modeling and predicting forest dynamics and their impacts on ecosystem services. In S. Das et al. (Eds.), *Application of Geospatial Technology and Modelling on Natural Resources Management. Advances in Geographic Information Science*. Springer, Cham. https://doi.org/10.1007/978-3-031-84698-4_7
- Yasin, E. H. E.**, Czimber, K., Gadallah, N. A. H., Mohammed, S. M. F., & Yagoub, Y. E. (2025). Use of geospatial technology and modeling to assess and predict climate change impacts. In S. C. Pal, U. Chatterjee, A. Saha, & D. Ruidas (Eds.), *Climate Change: Conflict and Resilience in the Age of Anthropocene. Advances in Global Change Research (Vol. 80)*. Springer, Cham. https://doi.org/10.1007/978-3-031-85359-3_6

- Yasin, E. H. E., Siddig, A. A. H., & Kornel, C. (2024).** Forests at the crossroads: Biodiversity conservation in the era of climate change. In *Environmental Sciences*. IntechOpen. <https://doi.org/10.5772/intechopen.1004224>
- Yasin, E. H. E., & Kornel, C. (2024).** Evaluating satellite image classification: Exploring methods and techniques. In *Geographic Information Systems - Data Science Approach*. IntechOpen. <https://doi.org/10.5772/intechopen.1003196>
- Yasin, H. E. E., Kornel, C., & Hemida, M. (2024).** Assessment and mapping of forest cover change in dryland, Sudan using remote sensing. In *Environmental Sciences*. IntechOpen. <https://doi.org/10.5772/intechopen.113862>
- Yasin, E. H. E., Siddig, A. A., Deiab, E. E., Czimer, K., Hasoba, A., & Osman, A. (2023).** Forest degradation in dryland ecosystems of Sudan: Review of the causes, consequences, assessment methods, and potential solutions. In *Environmental Sciences*. IntechOpen. <https://doi.org/10.5772/intechopen.113222>
- Osman, A. A. A., & Yasin, E. H. E. (2025).** Forest carbon stock and management strategies for riverine forests in Sudan. In *Forest Science: Advances Towards Sustainable Development and Climate Resilience*. IntechOpen. <https://doi.org/10.5772/intechopen.1010986>
- Younis, F. A., & Yasin, E. H. E. (2025).** The characteristics of popular hardwood species in Sudan—Review. In *Forest Science: Advances Towards Sustainable Development and Climate Resilience*. IntechOpen. <https://doi.org/10.5772/intechopen.1010985>

Conference papers

- Yasin, E. H. E. (2022).** Assessment of wood plant species composition, diversity, and structure analysis of Tozi Reserved Forest in Sinnar State, Sudan. In *Erdészeti Tudományok Konferencia, Sopron – 2022*. University of Sopron Press, pp. 91–101.
- Khiry, M. A., & Yasin, E. H. E. (2022).** GHG estimation from forest fire: A comparison of forest burned area using MODIS 64A1 and Sentinel-2 satellite images, savannah dryland region, Sudan. In *3rd International Marmara Scientific Research and Innovation Congress*. Iksad Publications, pp. 895–904.
- Albasha, A. K. O., & Yasin, E. H. E. (2022).** Impact of diluted Red Sea water on growth of Arak (*Salvadora persica*). In *Ankara International Congress on Scientific Research-VII*. Iksad Publications, pp. 1045–1056.
- Yasin, E. H. E., Eisawi, K. A. E., & Kornel, C. (2022).** Trees species composition, diversity, forest stand structure and regeneration status in Tozi Reserved Forest, Sinnar State, Sudan. In *3rd International Marmara Scientific Research and Innovation Congress*. Iksad Publications, pp. 905–919.